

The Expected Value Equal One Case Vance Harwood 19-Oct-2021

Consider the case of n independent, and identical random (iid) variables X_1, X_2, \dots, X_n , where the random variables X_i are lognormally distributed with parameters u_g and σ_g that satisfy the equation

$$u_g = -\frac{\sigma_g^2}{2}.$$

Where

- u_g = geometric mean, the average of $\ln(X_i)$
- σ_g = geometric standard deviation, the standard deviation of $\ln(X_i)$

What is the expected value $E[X_1 X_2 \dots X_n]$ as $n \rightarrow \text{Infinity}$?

The Status Quo

The status quo solution is based on the assertion that if random variables X_1 through X_n are independent and identically distributed, then the theorem below can be used. It asserts for independent random variables

$$E[X_1 X_2 \dots X_n] = E[X_i]^n \quad \text{Theorem 1} \quad (\text{MIT Random Variables and Expectation Corollary 10.2 page 31})$$

For a log normal distribution the expected value of $E[X_i] = e^{u_g + \frac{\sigma_g^2}{2}}$, substituting $-\frac{\sigma_g^2}{2}$ for u_g we have $e^{-\frac{\sigma_g^2}{2} + \frac{\sigma_g^2}{2}} = e^0 = 1$

According to theorem 1, $E[X_1 X_2 \dots X_n] = E[X_i]^n$ so $E[X_1 X_2 \dots X_n] = 1^n$ and $\lim_{n \rightarrow \infty} (1)^n = 1$.

Another Way

However, instead of using theorem 1 we can compute $E[X_1 X_2 \dots X_n]$ as $n \rightarrow \text{Infinity}$ directly.

The random variable for a log normal distribution is $X_i = e^{u_g + \sigma_g Z_i}$,

where Z_i = standard normal distribution $N(0,1)$.

Multiplying the random variables $X_1 X_2 \dots X_n$ we have $X_1 X_2 \dots X_n = (e^{u_g + \sigma_g Z_1})(e^{u_g + \sigma_g Z_2}) \dots (e^{u_g + \sigma_g Z_n})$. We can rewrite this as

$e^{n u_g} e^{n \sigma_g (Z_1 + Z_2 + \dots + Z_n)}$ which we can rewrite as

$$e^{n u_g} e^{n^2 \sigma_g \frac{(Z_1 + Z_2 + \dots + Z_n)}{n}}$$
 which equals $e^{n u_g} \left(e^{\sigma_g \frac{(Z_1 + Z_2 + \dots + Z_n)}{n}} \right)^{n^2}$.

For this case $\frac{\sigma_g^2}{2}$ must be positive therefore u_g must be negative, so as $n \rightarrow \infty$ the $\lim_{n \rightarrow \infty} e^{n u_g} = e^{-\infty}$ which equals zero.

The $\left(e^{\sigma_g \left(\frac{(Z_1+Z_2+\dots+Z_n)}{n} \right)} \right)^{n^2}$ term includes the expression $\left(\frac{(Z_1+Z_2+\dots+Z_n)}{n} \right)$ which by the law of large numbers (LLN) $\frac{(Z_1+Z_2+\dots+Z_n)}{n} \xrightarrow{a.s.} E[Z_i]$ as $n \rightarrow \infty$. Since $\lim_{n \rightarrow \infty} \left(e^{\sigma_g \left(\frac{(Z_1+Z_2+\dots+Z_n)}{n} \right)} \right)^{n^2} = \left[\lim_{n \rightarrow \infty} e^{\sigma_g \left(\frac{(Z_1+Z_2+\dots+Z_n)}{n} \right)} \right]^{n^2}$ we can invoke the LLN to say $= \left[\lim_{n \rightarrow \infty} e^{\sigma_g \left(\frac{(Z_1+Z_2+\dots+Z_n)}{n} \right)} \right]^{n^2} = \left[\lim_{n \rightarrow \infty} e^{\sigma_g E[Z_i]} \right]^{n^2}$. In this case $E[Z_i] = 0$, So we have $\left[\lim_{n \rightarrow \infty} e^0 \right]^{n^2} = (1)^{n^2} = 1$.

So the overall $e^{nu_g \left(e^{\sigma_g \left(\frac{(Z_1+Z_2+\dots+Z_n)}{n} \right)} \right)^{n^2}}$ expression, the product of the random variables reduces to $(0)(1) = 0$ as $n \rightarrow \infty$.

So as $n \rightarrow$ Infinity a direct calculation gives us $E[X_1 X_2 \dots X_n] = E[0] = 0$ for this case, which contradicts the solution based on theorem one.

The Coin Flip Contradiction Case Vance Harwood 19-Oct-2021

Take a fair coin, label the sides 1.2499 for heads and 0.7999 for tails. The expected value of the coin toss, which we will designate as the random variable X_i , is the arithmetic average of the sides, $\frac{(1.2499+0.7999)}{2}$, so $E[X_i] = 1.0249$. The product of a head/tail pair, $(1.2499)(0.7999)$ is approximately 0.9998. For a large number of tosses, what is the expected value of multiplying together the values of n coin tosses?

The Status Quo

Given that random variables X_1 through X_n are independent and identically distributed, the standard solution to this problem is to use Theorem 1 below. It asserts:

$$E[X_1 X_2 \dots X_n] = E[X_i]^n \quad \text{Theorem 1}$$

Since in this case $E[X_i] = 1.0249$, then according to Theorem 1 the expected value of the multiplied random variable $E[X_1 X_2 \dots X_n] = (1.0249)^n$. Since the $\lim_{n \rightarrow \infty} (1.0249)^n = \infty$ then $E[X_1 X_2 \dots X_n] = \infty$ also.

The Contradiction

However, consider the following calculation of $E[X_1 X_2 \dots X_n]$. There are three categories to consider as $n \rightarrow$ infinity:

1. The number of tails is larger than the number of heads
2. The number of heads equals the number of tails.
3. The number of heads is larger than the number of tails

The analysis for the first two categories can be combined and represents approximately half the outcomes of this exercise. If the number of tails in the $X_1 X_2 \dots X_n$ product is equal or greater than the number of heads then there are $\frac{n-(T-H)}{2}$ pairs of $(1.2499)(0.7999)$ in the product. Multiplying this term by $(0.7999)^{T-H}$ captures the contribution of any unpaired tails. The resulting equation is:

$$X_1 X_2 \dots X_n = ((1.2499)(0.7999))^{\frac{n-(T-H)}{2}} (0.7999)^{T-H}$$

In this scenario the absolute values of both of the base terms of the equation are positive numbers, less than one and the exponents are non-negative. Therefore, the overall product will always be less than one. In addition, as $n \rightarrow \infty$ for $y > 0$ and $y < 1$ the $\lim_{n \rightarrow \infty} (y)^n = 0$. So, we can say in the limit:

$$E[X_1 X_2 \dots X_n] = E\left[\lim_{n \rightarrow \infty} \left((0.7999)(1.2499)^{\frac{n-(T-H)}{2}} (0.7999)^{T-H} \right) \right] = E[0] = 0.$$

Since this is the outcome of the coin flip exercise approximately half the time, this result falsifies the status quo position that $E[X_1 X_2 \dots X_n] = (1.2499)^n$, where $\lim_{n \rightarrow \infty} (1.0249)^n \rightarrow \infty$.

The analysis of case 3, when the number of heads is larger than the number of tails also contradicts the status quo is below.

The Coin Flip Contradiction: More Heads Than Tails

The mean or expected value of the number of heads for each toss is 0.5 heads. If the coin toss is repeated n times, then as $n \rightarrow$ infinity the Law of Large Numbers says that the proportion of heads after n flips will almost surely converge to $\frac{1}{2}$ as $n \rightarrow$ infinity. While the absolute difference between the number of heads and tails can be large, the ratio of that difference divided by the total number of flips will go to zero as $n \rightarrow$ infinity, $\lim_{n \rightarrow \infty} \frac{|H-T|}{n} = 0$.

In the case of more heads than tails we have $\frac{n-(H-T)}{2}$ pairs of $(1.2499)(0.7999)$ multiplied by $(1.2499)^{H-T}$ which gives us the following equation:

$$E[X_1 X_2 \dots X_n] = E\left[\left((1.2499)(0.7999)\right)^{\frac{n-(H-T)}{2}} (1.2499)^{H-T}\right]$$

Rearranging the terms we have

$$\left((1.2499)(0.7999)\right)^{\frac{n-(H-T)}{2}} (1.2499)^{H-T} = \left((1.2499)(0.7999)\right)^{\frac{n}{2}} \left((1.2499)(0.7999)\right)^{-\frac{(H-T)}{2}} (1.2499)^{H-T} .$$

Taking the terms to the $\frac{n}{2}$ power and rearranging we have:

$$\left(\left((1.2499)(0.7999)\right)^{\frac{1}{2}} \left((1.2499)(0.7999)\right)^{-\frac{1}{2} \left(\frac{H-T}{n}\right)} (1.2499)^{\left(\frac{H-T}{n}\right)}\right)^n .$$

$$\text{The } \lim_{n \rightarrow \infty} \left(\left((1.2499)(0.7999)\right)^{\frac{1}{2}} \left((1.2499)(0.7999)\right)^{-\frac{1}{2} \left(\frac{H-T}{n}\right)} (1.2499)^{\left(\frac{H-T}{n}\right)}\right)^n = \left[\lim_{n \rightarrow \infty} \left(\left((1.2499)(0.7999)\right)^{\frac{1}{2}} \left((1.2499)(0.7999)\right)^{-\frac{1}{2} \left(\frac{H-T}{n}\right)} (1.2499)^{\left(\frac{H-T}{n}\right)}\right)\right]^n .$$

Since the limit of the absolute difference between the number of heads and tails divided by the total number of flips $\lim_{n \rightarrow \infty} \frac{|H-T|}{n} = 0$, then as $n \rightarrow$ infinity the $\frac{(H-T)}{n}$ terms go to zero giving us

$$\lim_{n \rightarrow \infty} \left((1.2499)(0.7999)\right)^{-\frac{1}{2} \left(\frac{H-T}{n}\right)} = 1 \text{ and } \lim_{n \rightarrow \infty} (1.2499)^{\left(\frac{H-T}{n}\right)} = 1 \text{ so the overall limit as } n \rightarrow \infty \text{ is}$$

$$\lim_{n \rightarrow \infty} \left(\left((1.2499)(0.7999)\right)^{\frac{1}{2}} (1)(1)\right) = \left((1.2499)(0.7999)\right)^{\frac{1}{2}} . \text{ Since } \left((1.2499)(0.7999)\right)^{\frac{1}{2}} \text{ is less than one}$$

and greater than zero then $\lim_{n \rightarrow \infty} \left(\left((1.2499)(0.7999)\right)^{\frac{1}{2}}\right)^n = 0$. Therefore for the heads greater than tails

$$\text{case } E[X_1 X_2 \dots X_n] = E\left[\lim_{n \rightarrow \infty} \left(\left((1.2499)(0.7999)\right)^{\frac{1}{2}} \left((1.2499)(0.7999)\right)^{-\frac{1}{2} \left(\frac{H-T}{n}\right)} (1.2499)^{\left(\frac{H-T}{n}\right)}\right)^n\right] = E[0] = 0$$

which also contradicts the status quo result of $E[X_1 X_2 \dots X_n] = (1.2499)^n$, where $\lim_{n \rightarrow \infty} (1.2499)^n \rightarrow \infty$

Two Means Straddling One Case

Vance Harwood 19-Oct-2021

The Case

Consider the case of n independent, and identical random (iid) variables X_1, X_2, \dots, X_n , where the random variables X_i take on positive real number values of b or c with 0.5 probability. The characteristics of the values b and c are:

1. v is a positive, non-infinite real number
2. d is a negative, non-infinite real number
3. $b = e^{(d+v)}$ and $c = e^{(d-v)}$
4. $e^d > \frac{2}{(e^v + e^{-v})}$

Example parameters that meet these conditions are $v = 0.2$ and $d = -0.01$.

The net effects of these restrictions are:

1. That b multiplied by $c = e^{(d+v+d-v)} = e^{(2d)}$
2. The arithmetic mean of b and c will always be greater than one
3. The geometric mean of b and c will always be less than one
4. The random variable $E[X_i]$ has a defined mean and variance

The Status Quo

The expected value of $E[X_i]$ is the arithmetic mean of X_i , which is

$$\frac{e^{(d+v)} + e^{(d-v)}}{2} \text{ which can be rewritten as } e^d \left(\frac{e^{(v)} + e^{(-v)}}{2} \right)$$

The term on the left, e^d was defined earlier as being more positive than $\frac{2}{(e^v + e^{-v})}$. The term $\frac{2}{(e^v + e^{-v})}$ is the reciprocal of the right-hand term $\frac{e^{(v)} + e^{(-v)}}{2}$. A number multiplied by its reciprocal equals one, but since e^d is defined to be more positive than $\frac{2}{(e^v + e^{-v})}$ the product of e^d and $\left(\frac{e^{(v)} + e^{(-v)}}{2} \right)$ must be larger than one. Therefore, the expected value, $E[X_i]$ must be greater than 1.

Given that, theorem 1 states that in this case, $E[X_1 X_2 \dots X_n] = E[X_i]^n$ which goes to ∞ as $n \rightarrow \infty$

The Contradiction

$E[X_1 X_2 \dots X_n]$ is equal to $E[e^{\ln(X_1 X_2 \dots X_n)}]$ we can further rewrite this as

$E[e^{(\ln(X_1) + \ln(X_2) + \dots + \ln(X_n))}]$. We multiply the exponent by $\frac{n}{n}$ which we rewrite as

$$\mathbb{E}\left[e^{n\left(\frac{1}{n}(\ln(X_1)+\ln(X_2)+\dots+\ln(X_n))\right)}\right]. \quad \text{Equation (1)}$$

Notice that the term $\frac{1}{n}(\ln(X_1) + \ln(X_2) + \dots + \ln(X_n))$ is the sample average of all the $\ln(X_i)$ terms

By the Law of Large Numbers we can say the sample average converges almost surely to the expected value $\mathbb{E}[\ln(X_i)]$ as $n \rightarrow \infty$.

The natural log of the two values b and c of X_i are $\ln(e^{(d+v)})$ and $\ln(e^{(d-v)})$ respectively, which reduces to $d+v$ and $d-v$. The expected value then of $\mathbb{E}[\ln(X_i)]$ is calculated as $(d+v)*0.5$ plus $(d-v)*0.5 = 0.5(d+v+d-v) = 0.5(2d) = d$.

Returning to equation 1 we can now say that

$$\begin{aligned} \mathbb{E}\left[e^{n\left(\frac{1}{n}(\ln(X_1)+\ln(X_2)+\dots+\ln(X_n))\right)}\right] &= \mathbb{E}\left[\left(e^{\left(\frac{1}{n}(\ln(X_1)+\ln(X_2)+\dots+\ln(X_n))\right)}\right)^n\right] \text{ Since} \\ \lim_{n \rightarrow \infty} \left[\left(e^{\left(\frac{1}{n}(\ln(X_1)+\ln(X_2)+\dots+\ln(X_n))\right)}\right)^n\right] &= \left[\lim_{n \rightarrow \infty} \left(e^{\left(\frac{1}{n}(\ln(X_1)+\ln(X_2)+\dots+\ln(X_n))\right)}\right)\right]^n \text{ then by LLN} \\ \left[\lim_{n \rightarrow \infty} \left(e^{\left(\frac{1}{n}(\ln(X_1)+\ln(X_2)+\dots+\ln(X_n))\right)}\right)\right]^n &\xrightarrow{a.s.} \left[e^{\mathbb{E}[\ln(X_i)]}\right]^n. \text{ Since } \mathbb{E}[\ln(X_i)] = d \text{ we have} \end{aligned}$$

$$\left[e^{\mathbb{E}[\ln(X_i)]}\right]^n = \left[e^d\right]^n \text{ and since } d \text{ is always negative then } e^d < 1 \text{ and the } \lim_{n \rightarrow \infty} (e^d)^n = 0$$

Therefore in this case $\mathbb{E}[X_1 X_2 \dots X_n] = \mathbb{E}[0] = 0$ which contradicts theorem 1's assertion that $\mathbb{E}[X_1 X_2 \dots X_n]$ will go to infinity as $n \rightarrow \infty$